Choose the Right Car Seat based on a child's weight, height, age and development.

Infant Car Seats

Protect a baby in a crash.





Convertible Car Seats

Protect a child from birth to 5 or 6 years old.



Booster Seats

Position the seat belt correctly on kids less than 4'9" tall or younger than 8 years old.



Choose the Right Car Seat

based on a child's weight, height, age and development.

Infant Car Seat

- Rear-facing only.
- Weight and height limits vary by the car seat.
- Recline seat between 30-45° according to level indicator or instructions.
- Use a convertible car seat when a baby reaches the weight or height limit on the infant car seat.

Convertible Car Seat

- Rear-facing then transitions to forward-facing.
- Most come with an infant insert and can be used for rear-facing at birth.
- Children stay rear-facing until they reach the rear-facing weight or height limit for the car seat.
- Once the child reaches the limit and is at least 2 years old, he/she begins forward-facing.
- Children should stay forward-facing in the 5-point harness until reaching the weight or height limit for the car seat.
- When forward-facing, attach tether located on the top of the car seat to correct tether anchor in the vehicle.

Booster Seat

- Helps position a seat belt properly on a child.
- Incorrect seat belt fit can cause abdominal and neck injuries or ejection in a crash.
- Ohio Booster Seat Law: Children must be at least 8 years old to stop using a booster seat unless they are at least 4 feet 9 inches tall.

Remember:

- Always buckle a child into the car seat.
- Rear facing is safer.
- To find a car seat fitting station near you, visit: www.nhtsa.gov/equipment/car-seats-and-booster-seats#install-inspection.



Original content of this guide was developed by Columbus Public Health's Safe Kids Central Ohio program and made possible with support from State Farm; 12/2017.

This publication (Exam Room Card) was supported by the Cooperative Agreement number, 6 NB010T009137-01-02, funded by the Centers for Disease control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services.